# Contents

Preface vii

Editorial Note x

**I. Composite fermions and fractional quantum Hall effect**

Resonant tunneling in the quantum Hall regime: measurement of fractional charge
V.J. Goldman 1

Recent progress in the theory of composite fermions near even-denominator filling factors
N. Read 7

Composite fermion theory, edge currents, and the fractional quantum Hall effect
G. Kirczenow and B.L. Johnson 13

The Aharonov–Bohm effect in the fractional quantum Hall regime
J.D.F. Franklin, I. Zailer, C.J.B. Ford, P.J. Simpson, J.E.F. Frost, D.A. Ritchie, M.Y. Simmons and M. Pepper 17

The unifying role of effective field in the composite fermion model of the fractional quantum Hall effect
D.R. Leadley, M. van der Burgt, R.J. Nicholas, P.J. Gee, J. Singleton, J.J. Harris and C.T. Foxon 22

Composite fermions around Landau level filling factor $\nu = 3/2$
R.R. Du, A.S. Yeh, H.L. Stormer, D.C. Tsui, L.N. Pfeiffer and K.W. West 26

Magnetoexciton luminescence at 1/3 Landau filling as a function of electron–hole separation
F. Plentz, D. Heiman, A. Pinczuk, L.N. Pfeiffer and K.W. West 30

Time-resolved phonon absorption in the fractional quantum Hall regime
J.E. Digby, U. Zeitler, C.J. Mellor, A.J. Kent, K.A. Benedict, L.J. Challis, J.R. Middleton and T. Cheng 34

Composite fermions examined with surface acoustic waves
R.L. Willett and L.N. Pfeiffer 38

The physical significance of singularities in the Chern–Simons Fermi liquid description of a partially-filled Landau level
A. Stern and B.I. Halperin 42

Universal behaviour of the thermoelectric power of composite fermions
B. Tieke, U. Zeitler, R. Fletcher, S.A.J. Wiegers, A.K. Geim, J.C. Maan and M. Henini 46

Evidence for composite fermions from the magnetothermopower of 2D holes
P.A. Crump, B. Tieke, R.J. Barraclough, B.L. Gallagher, R. Fletcher, J.C. Maan, T.M. Fromhold and M. Henini 50

Electron spin resonance in AlGaAs/GaAs in the regime of fractional filling
R. Meisels, I. Kulac, G. Sundaram, F. Kuchar, B.D. McCombe, G. Weimann and W. Schlapp 55

Mesoscopic transport properties of composite fermions
A.S. Sachrajda, Y. Feng, H.A. Carmona, A.K. Geim, P.C. Main, L. Eaves and C.T. Foxon 59

Transport properties of composite fermions in narrow cross junctions
J. Herfort, Y. Takagaki, R. Hey, K.J. Friedland, K. Ploog, D.K. Maude, J.C. Portal, J. Takahara and K. Gamo 63

Ballistic conductance of composite fermions
A. Khaetskii, Yu.V. Nazarov and G.E.W. Bauer 67

Measurements on a composite fermion split-gate
C.-T. Liang, C.G. Smith, D.R. Mace, J.T. Nicholls, J.E.F. Frost, M.Y. Simmons, A.R. Hamilton, D.A. Ritchie and M. Pepper 71

Critical behavior in the fractional quantum Hall effect
B. Huckestein 75

Understanding the dynamics of fractional edge states with composite fermions
D.B. Chklovskii and B.I. Halperin 79

Composite-fermion picture for the double-layer fractional quantum Hall effect
T. Nakajima and H. Aoki 83

Resonant Raman scattering and numerical study in the fractional quantum Hall regime
S. He and P.M. Platzman 87

Effective masses of composite fermions
N. d'Ambrumenil and R. Morf 92

Breakdown phenomena of integer and fractional quantum Hall effects under pulsed magnetic fields
T. Takamasu, H. Dodo, M. Ohno and N. Miura 95

Influence of scattering processes on the electronic properties of composite fermions
I.V. Kukushkin, R.J. Haug, K. von Klitzing and K. Eberl 99

Pulse propagation in spin-polarized edge channels and at fractional filling factors
G. Ernst, N.B. Zhitenev, R.J. Haug, K. von Klitzing and K. Eberl 102

**II. Parallel quasi-2D layers**

Correlated insulating states of an interacting bilayer electron system
H.C. Manoharan, Y.W. Suen, M.B. Santos and M. Shayegan 106

The thermodynamics of a charged fermion layer at high $r_s$ values
S. Shapira, U. Sivan, P.M. Solomon, E. Buchstab, M. Tischler and G. Ben Yoseph 113

Observation of correlated $\nu = 1$ quantum Hall and insulating states in strongly coupled p-type double quantum wells
R.J. Hyndman, T. Ihn, B.L. Gallagher, P.J. Rodgers, M. Henini, T.J. Foster and J.R. Middleton 117

Josephson phenomena in bilayer quantum Hall systems
Z.F. Ezawa 122

Single particle and electron–electron scattering rates in coupled quantum wells
Y. Berk, A. Kamenev, A. Palevski, H. Shtrikman and M. Slutzky 126

Strong enhancement of Coulomb drag in double-layer systems due to correlations among carriers
L. Świerkowski, J. Szymański and Z.W. Gortel 130

Frictional drag between closely spaced two-dimensional electron gases
H. Rubel, E.H. Linfield, N.P.R. Hill, J.T. Nicholls, D.A. Ritchie, K.M. Brown, M. Pepper and G.A.C. Jones 134

Tuning a double quantum well Fermi surface with in-plane magnetic fields
J.A. Simmons, N.E. Harff, T.M. Eiles, S.K. Lyo and J.F. Klem 138

Suppression of resonant tunneling in a coupled quantum well
Y. Ohno and H. Sakaki 142

Superfluid ground state in electron–hole double layer systems
Z.W. Gortel and L. Świerkowski 146

An unusual superconductivity mechanism in systems with pairing of spatially separated electrons and holes
S.I. Shevchenko 150

Coulomb blockade as a non-invasive probe in double layer 2DEG systems
M. Field, C.G. Smith, M. Pepper, K.M. Brown, E.H. Linfield, M.P. Grimshaw, D.A. Ritchie and G.A.C. Jones 154

Observation of many-body interactions of electrons in coupled double quantum wells
A.S. Plaut, A. Pinczuk, B.S. Dennis, J.P. Eisenstein, L.N. Pfeiffer and K.W. West 158

Stability and dynamical properties of a double-layer Wigner crystal in two dimensions
G. Goldoni, V. Schweigert and F.M. Peeters 163

Tunneling between parallel two-dimensional electron liquids
T. Jungwirth and A.H. MacDonald 167

Magnetotransport of a low-disorder triple-layer electron system in perpendicular or parallel magnetic fields
T.S. Lay, S.P. Shukla, J. Jo, X. Ying and M. Shayegan 171

**III. Resonant tunnelling and tunnelling**

Multiphoton-assisted tunneling, dynamic localization and absolute negative conductance
B.J. Keay, S.J. Allen, Jr., K.D. Maranowski, A.C. Gossard, U. Bhattacharya and M.J.W. Rodwell 176

Non-equilibrium acoustic phonon-assisted tunnelling in GaAs/(AlGa)As double barrier devices
F.F. Ouali, N.N. Zinov'ev, L.J. Challis, A.V. Akimov, D.N. Hill, F.W. Sheard and M. Henini 181

Field enhanced blockade of the confined energy levels in nanometer scale pillar arrays
B.W. Alphenaar, Z.A.K. Durani, M. Wagner and K. Köhler 185

Spin–orbit splitting of the spectrum of 2D electrons in the surface quantum well of gapless semiconductors
G.M. Minkov, A.V. Germanenko, V.A. Larionova and O.E. Rut 189

Landau assisted vertical transport in MQW p–i–n GaAs/AlGaAs diodes
R.J. Stone, J.G. Michels, S.L. Wong, C.T. Foxon, R.J. Nicholas and A.M. Fox 192

An optical and electrical study of tunnelling mechanisms through indirect gap single barrier GaAs/AlAs/GaAs heterostructures
J.J. Finley, R. Teissier, M.S. Skolnick, J.W. Cockburn, R. Grey, G. Hill and M.A. Pate 197

Interband tunneling of holes through single-barrier nanostructures
M. Morifuji, K. Taniguchi and C. Hamaguchi 201

Interface composition dependence of the band offset in InAs/GaSb
M.S. Daly, D.M. Symons, M. Lakrimi, R.J. Nicholas, N.J. Mason and P.J. Walker 205

Transfer efficiency of hot electrons in a metal($CoSi_2$)/insulator($CaF_2$) quantum interference transistor
T. Suemasu, W. Saitoh, Y. Kohno, K. Mori, M. Watanabe and M. Asada 209

Resonant tunneling through a pseudo-quantum well in single barrier heterostructure
Yu.V. Dubrovskii, Yu.N. Khanin, E.E. Vvdovin, I.A. Larkin and T.G. Andersson 213

Resonant tunneling in time-dependent fields through laterally confined double barriers
R. Aguado, J. Iñarrea and G. Platero 217

Broadening of resonance level induced by interface roughness scattering in double barrier tunneling structures
T. Hoshida, M. Tsuchiya and T. Kamiya 222

Charge accumulation over a region of electrical multistability in a double barrier structure
M.L.F. Lerch, D.J. Fisher, A.D. Martin, C. Zhang and L. Eaves 226

Origin of current broadening in resonant tunneling via localised states
C. Zhang, D.J. Fisher and S.M. Stewart 231

Effect of spacer layer on quantum interference in double-barrier resonant tunneling structures
A.E. Belyaev, S.A. Vitusevich, T. Figielski, B.A. Glavin, R.V. Konakova, L.N. Kravchenko, A. Makosa and T. Wosinski 235

Ultra-high-frequency coherent sound generation in resonant tunneling
S.S. Makler, D.E. Tuyarot, E.V. Anda and M.I. Vasilevskiy 239

Anomalous transport of indirect excitons in coupled AlAs/GaAs quantum wells
L.V. Butov, A. Zrenner, M. Hagn, G. Abstreiter, G. Böhm and G. Weimann 243

Resonant tunnelling through $D^-$ states
J.G.S. Lok, A.K. Geim, J.C. Maan, I. Marmorkos, F.M. Peeters, N. Mori, L. Eaves, P. McDonnell, M. Henini and P.C. Main 247

Resonant tunneling through two impurities in disordered barriers
A.K. Savchenko, V.V. Kuznetsov, A. Woolfe, D.R. Mace, M. Pepper, D.A. Ritchie and G.A.C. Jones 251

**IV. Integer quantum Hall effect**

Activated conductivity in the quantum Hall effect
M.M. Fogler, D.G. Polyakov and B.I. Shklovskii 255

NMR measurement of the spin magnetization and spin dynamics in the quantum Hall regimes
S.E. Barrett, G. Dabbagh, L.N. Pfeiffer, K.W. West and R. Tycko 261

Imaging of edge channels in the integer quantum Hall regime by the lateral photoelectric effect
R.J.F. van Haren, F.A.P. Blom, W. de Lange and J.H. Wolter 267

Voltage distribution and phase-breaking scattering in the quantum Hall regime
T. Ando 270

The 2D electron gas near $\nu = 1$ as a Skyrme crystal
L. Brey, H.A. Fertig, R. Côté and A.H. MacDonald 274

Collapse of the quantum Hall state by floating-up of extended-state bands with increasing disorder
T. Okamoto, Y. Shinohara, S. Kawaji and A. Yagi 278

*I–V* characteristic of a 2D electron channel with the filling factor close to integer
V.B. Shikin 282

Quantum Hall effect with strong spin–orbit mixing of Zeeman states
D.G. Polyakov and M.E. Raikh 285

Potential drops across quantum Hall effect samples – in the bulk or near the edges?
W. Dietsche, K. von Klitzing and K. Ploog 289

**V. Hybrid superconductor– and magnet–semiconductor systems**

Supercurrent and quasi-particle transport in a two-dimensional electron gas with superconducting electrodes
B.J. van Wees 293

Observation of the quantized critical current in a superconducting quantum point contact
H. Takayanagi, T. Akazaki and J. Nitta 298

Theory of Andreev resonances in quantum dots
M. Leadbeater, N.R. Claughton, C.J. Lambert and V.N. Prigodin 302

Superconductivity and the Josephson effect in a periodic array of Nb–InAs–Nb junctions
H. Drexler, J.G.E. Harris, E.L. Yuh, K.C. Wong, S.J. Allen, E.G. Gwinn, H. Kroemer and E.L. Hu 306

Size-scaling experiment in a two-dimensional electron gas using inhomogeneous magnetic field due to a superconducting gate
A.K. Geim, S.V. Dubonos, I.V. Grigorieva and S.J. Bending 311

Far-infrared studies of induced superconductivity in quantum wells
E.L. Yuh, J.G.E. Harris, T. Eckhause, K.C. Wong, E. Gwinn, H. Kroemer and S.J. Allen 315

Transport through a 2DEG channel with superconducting boundaries
J. Nitta, B.J. van Wees, J.P. Heida, T.M. Klapwijk, A. Dimoulas, W. van de Graaf and G. Borghs 320

Transport through superconductor–semiconductor junctions in different scattering limits
D.A. Williams, A.M. Marsh and H. Ahmed 324

Two-dimensional electrons in a lateral magnetic superlattice
H.A. Carmona, A. Nogaret, A.K. Geim, P.C. Main, T.J. Foster, M. Henini, S.P. Beaumont, H. McLelland and M.G. Blamire 328

Magnetoresistance oscillation in a two-dimensional electron gas under periodic modulation of electric and magnetic fields
A. Endo, S. Izawa, S. Katsumoto and Y. Iye 333

Magnetotransport in periodic magnetic fields
P.D. Ye, D. Weiss, R.R. Gerhardts, K. von Klitzing, K. Eberl and H. Nickel 337

Electrons in a periodic magnetic field
I.S. Ibrahim and F.M. Peeters 341

Two-dimensional electron gas in a spatially varying magnetic field
E. Hofstetter, J.M.C. Taylor and A. MacKinnon 345

Magnetoresistance of a two-dimensional electron gas in random magnetic fields
L. Theil Hansen, R. Taboryski, A. Smith, P.E. Lindelof and P. Hedegård 349

Microwave-induced "Somersault effect" in the flow of Josephson current through a quantum constriction
L.Y. Gorelik, V.S. Shumeiko, R.I. Shekhter, G. Wendin and M. Jonson 353

**VI. Optical studies of quantum wells and superlattices**

Optical spectroscopy of neutral and charged excitons in GaAs/AlGaAs quantum wells in high magnetic fields
G. Finkelstein, H. Shtrikman and I. Bar-Joseph 357

Negatively charged excitons ($X^-$) and $D^-$ triplet transitions in GaAs/$Al_{0.3}Ga_{0.7}As$ multiple quantum wells
S.R. Ryu, W.-Y. Yu, L.P. Fu, Z.X. Jiang, A. Petrou, B.D. McCombe and W. Schaff 363

Coherent submillimeter-wave emission from non-equilibrium two-dimensional free carrier plasmas in AlGaAs/GaAs heterojunctions
K. Hirakawa, I. Wilke, K. Yamanaka, H.G. Roskos, M. Voßebürger, F. Wolter, C. Waschke, H. Kurz, M. Grayson and D.C. Tsui 368

Magnetic field dependence of many-body processes in quantum well luminescence
H.P. van der Meulen, J. Rubio, J.M. Calleja, H. Schweizer, F. Scholz, V. Härle and R. Bergmann 372

Phase of magnetic oscillation of luminescence energy in asymmetric quantum wells
T. Tsuchiya, S. Katayama and T. Ando 376

Strain-induced suppression of spin splitting in asymmetric p-type quantum wells
O. Mauritz and U. Ekenberg 380

Dependence of the optical absorption on the in-plane light polarization of QWs grown on low symmetry (011) and (113) surfaces
J. Los and A. Fasolino 384

Magnetooscillations in the electroluminescence intensity of excited subbands in resonantly coupled superlattices
D. Bertram, H.T. Grahn, O. Kuhn, D.K. Maude, J.C. Portal, K. Ploog and K. von Klitzing 387

Photoluminescence studies of current-induced non-equilibrium population in magnetically quantized two-dimensional electron gases
N.N. Zinov'ev, A.V. Andrianov, L.J. Challis, C.T. Foxon and J.J. Harris 392

Resonant bound bipolarons in a superlattice in a high magnetic field
J.M. Shi, F.M. Peeters and J.T. Devreese 397

Nonlinear resonant optical rectification in a coupled quantum well
K. Unterrainer, J.N. Heyman, K. Craig, B. Galdrikian, M.S. Sherwin, H. Drexler, K. Campman, P.F. Hopkins and A.C. Gossard 401

Cyclotron resonance in short period $(GaAs)_n/(AlAs)_n$ superlattices
T. Fukuda, K. Yamanaka, H. Momose, N. Mori, C. Hamaguchi, Y. Imanaka, Y. Shimamoto and N. Miura 406

Cyclotron resonance and subband-Landau level coupling in 2D electron and hole gases
R. Winkler 411

Cyclotron resonance in InAs quantum wells in tilted magnetic fields
Yu.B. Vasilyev, S.V. Ivanov, B.Ya. Meltser, S.D. Suchalkin and P. Grambow 415

Optical and transport studies of highly acceptor doped GaAs/AlGaAs quantum wells
A.C. Ferreira, A.V. Buyanov, P.O. Holtz, B.E. Sernelius, B. Monemar, O. Mauritz, U. Ekenberg, M. Sundaram, K. Campman, J.L. Merz and A.C. Gossard 420

Mobile excitons and Fermi edge singularities in an interacting 2D electron gas
J.A. Brum, S. Brown, P. Hawrylak, J.F. Young and Z. Wasilewski 424

Excitons in a dense two-dimensional electron gas at a semiconductor surface
P.D. Altukhov 428

High density, spatially separated electron/hole plasmas in mixed type I–type II GaAs/AlAs heterostructures
P.D. Buckle and P. Dawson 431

Investigation of *g*-factors, Zeeman splittings, exchange interactions and field-dependent spin relaxation in III–V quantum wells
N.J. Traynor, M.J. Snelling, R.T. Harley, R.J. Warburton and M. Hopkinson 435

The $D^-$ bound exciton observed in GaAs/AlGaAs quantum wells
P.O. Holtz, Q.X. Zhao, B. Monemar, M. Willander, K. Campman, M. Sundaram, J.L. Merz and A.C. Gossard 439

Trion–electron scattering in GaAs/AlAs quantum wells
A. Manassen, R. Harel, E. Cohen, A. Ron, E. Linder and L.N. Pfeiffer 443

Creation and annihilation of positively and negatively charged excitons in GaAs quantum wells
H. Buhmann, P.H. Beton, L. Eaves and M. Henini 447

Magneto-optical spectroscopy of neutral and negatively charged excitons in GaAs quantum wells
A.J. Shields, M. Pepper, M.Y. Simmons and D.A. Ritchie 451

Influence of the electron–electron and electron–phonon interaction on the cyclotron resonance of 2DEG in GaAs
C.M. Hu, E. Batke, K. Köhler and P. Ganser 456

Electron and hole *g*-factor oscillations in CdTe-based modulation-doped quantum wells
J.X. Shen, W. Ossau, F. Fischer, A. Waag and G. Landwehr 460

Effective mass anisotropy and many-body effects in 2D GaAs/(Ga, < ZW > Al)As hole gases observed in very high magnetic fields: comparison of theory and experiment
B.E. Cole, S.O. Hill, Y. Imanaka, Y. Shimamoto, W. Batty, J. Singleton, J.M. Chamberlain, N. Miura, M. Henini and T. Cheng 464

Photon drag saturation spectroscopy of a two-dimensional electron system
H. Sigg, P. van Son and K. Köhler 468

Zero-field spin-splitting in InAs/AlSb quantum wells
C. Gauer, M. Hartung, A. Wixforth, J.P. Kotthaus, B. Brar and H. Kroemer 472

Fermi-edge singularity of the Tomonaga–Luttinger liquids in a magnetic field
T. Ogawa and H. Otani 476

**VII. Lateral transport in 2D systems**

Observation of the zero Hall resistance state in a semimetallic superlattice
M.S. Daly, K.S.H. Dalton, M. Lakrimi, N.J. Mason, D.K. Maude, R.J. Nicholas, J.C. Portal, D.M. Symons, M. van der Burgt and P.J. Walker 481

Two-dimensional electron liquid in a weak magnetic field
L.I. Glazman and I.L. Aleiner 485

Phonon assisted energy relaxation for 2D electrons at high magnetic field
S.V. Iordanskii, Y. Levinson and S. Levit 489

Electron density dependence of the effective *g*-factor in a two-dimensional electron gas at strong magnetic fields
J.H. Oh, K.J. Chang, G. Ihm and S.J. Lee 493

Inefficiency of odd relaxation and propagation of electron beams in a two-dimensional electron system
R.N. Gurzhi, A.N. Kalinenko and A.I. Kopeliovich 497

Energy band diagrams of nano-size field-effect transistors
Y. Fu and M. Willander 500

The Hofstadter energy spectrum for an interacting 2DEG
V. Gudmundsson and R.R. Gerhardts 505

Fermi contour of the 2D electron layer at GaAs/$Al_xGa_{1-x}As$ heterojunction subject to in-plane magnetic fields
L. Smrčka, P. Vašek, J. Koláček, T. Jungwirth and M. Cukr 509

Coulomb interaction effects on the magnetoconductivity of laterally modulated two-dimensional electron systems
A. Manolescu, R.R. Gerhardts, M. Tornow, D. Weiss. K. von Klitzing and G. Weimann 513

Edge of the two-dimensional electron gas in a gated heterostructure
I.A. Larkin and J.H. Davies 517

$X_z$–$X_{xy}$ crossover in a two-dimensional electron gas in AlAs
A.F.W. van de Stadt, P.M. Koenraad, J.A.A.J. Perenboom and J.H. Wolter 521

Magnetotransport in wide parabolic PbTe quantum wells
J. Oswald, G. Heigl, M. Pippan, G. Span, T. Stellberger, D.K. Maude and J.C. Portal 525

Oscillations in the low-field diffusion thermopower of a 2D electron gas
R. Fletcher, P.T. Coleridge and Y. Feng 529

Magnetoresistance anomaly of a two-dimensional electron gas in tilted magnetic fields
P. Středa, P. Vašek and M. Cukr 533

Electron–phonon scattering rates in GaAs/AlGaAs 2DEG samples below 0.5 K
A. Mittal, R.G. Wheeler, M.W. Keller, D.E. Prober and R.N. Sacks 537

Tilted magnetic field studies of spin- and valley-splittings in Si/$Si_{1-x}Ge_x$ heterostructures
P. Weitz, R.J. Haug, K. von Klitzing and F. Schäffler 542

Negative magnetoresistance and electron–electron interaction in Si:SiGe quantum wells
T.J. Thornton, A. Matsumura and J. Fernández 547

Low temperature magnetotransport of 2D electron and hole systems in high-mobility Si–$Si_{1-x}Ge_x$ heteroctructures
R.B. Dunford, R. Newbury, V.A. Stadnik, F.F. Fang, R.G. Clark, R.H. McKenzie, R.P. Starrett, E.E. Mitchell, P.J. Wang, J.O. Chu, K.E. Ismail and B.S. Meyerson 550

Conduction band spin splitting and negative magnetoresistance in $A_3B_5$ heterostructures
G.E. Pikus and F.G. Pikus 556

Effective masses in high-mobility 2D electron gas structures
P.T. Coleridge, M. Hayne, P. Zawadzki and A.S. Sachrajda 560

Magnetoresistance oscillations due to intersubband scattering in a two-dimensional electron system
T.H. Sander, S.N. Holmes, J.J. Harris, D.K. Maude and J.C. Portal 564

Direct measurements of energy relaxation time of electrons in AlGaAs/GaAs heterostructures under quasi-equilibrium conditions
A.A. Verevkin, N.G. Ptitsina, G.M. Chulcova, G.N. Gol'tsman, E.M. Gershenzon and K.S. Yngvesson 569

An elegant verification of the negative charge-state of the *DX*-centre from mobility measurements of 2D electrons
M. Hayne, A. Usher, J.J. Harris and C.T. Foxon 574

Mobility and quantum lifetime in a GaAs/AlGaAs heterostructure: tuning of the remote-charge correlations
P. Wisniewski, T. Suski, E. Litwin-Staszewska, G. Brunthaler and K. Köhler 579

Anisotropy of the conductivity in δ-doped multilayers
Th. Ihn, C. Gauer, F. Koch and J.P. Kotthaus 583

Electron transport in a non-planar 2DEG
M.L. Leadbeater, C.L. Foden, J.H. Burroughes, T.M. Burke, L.L. Wang, M.P. Grimshaw, D.A. Ritchie and M. Pepper 587

**VIII. Quantum dots and wires – transport**

High-frequency transport through mesoscopic structures
L.P. Kouwenhoven, N.C. van der Vaart, Yu.V. Nazarov, S. Jauhar, D. Dixon, K. McCormick, J. Orenstein, P.L. McEuen, Y. Nagamune, J. Motohisa and H. Sakaki 591

Photon-assisted tunneling through a double quantum dot
R.H. Blick, R.J. Haug, K. von Klitzing and K. Eberl 595

Time-resolved single-electron tunnelling between Landau states in a quantum dot
N.C. van der Vaart, M.P. de Ruyter van Steveninck, F. Cochius, L .P. Kouwenhoven, A.T. Johnson, Yu.V. Nazarov, C. Harmans and C.T. Foxon 600

Electrophonon resonances and optical phonon generation in nanowires
V.B. Pevzner, V.L. Gurevich and G.J. Iafrate 605

Detection of the oscillation of the Fermi energy of a 2DEG
C.H.W. Barnes, D.R. Mace, G. Faini, D. Mailly, M.Y. Simmons, C.J.B. Ford and M. Pepper 608

Quantum box energy spectroscopy by 3D–0D resonant tunnelling
G. Faini, C. Vieu, F. Laruelle, P. Krauz, E. Bedel and C. Fontaine 613

Calculated *I–V* characteristics of low-dimensional structures
M. Boero and J.C. Inkson 618

Theory of Coulomb blockade of tunneling through a double quantum dot
K.A. Matveev, L.I. Glazman and H.U. Baranger 623

Tunnelling experiments through coupled zero-dimensional systems
A.S. Sachrajda, C. Gould, Y. Feng, M.W.C. Dharma-wardana and P.T. Coleridge 627

Coulomb gap of coupled quantum dots with adjustable interdot tunneling
C.H. Crouch, C. Livermore, F.R. Waugh, R.M. Westervelt, K.L. Campman and A.C. Gossard 631

Linear and non-linear transport through coupled quantum dots
D. Dixon, L.P. Kouwenhoven, P.L. McEuen, Y. Nagamune, J. Motohisa and H. Sakaki 636

Wave function and conductance calculations for a quantum dot at resonant tunnelling peaks
P.N. Butcher and J.A. McInnes 640

Magneto-tunnelling spectroscopy of a quantum dot charged with a few electrons
A. Nogaret, M.J. Gompertz, P.C. Main, L. Eaves, T.J. Foster, M. Henini and S.P. Beaumont 644

Ground states of 2D and 3D quantum dots
L.D. Hallam, N.A. Bruce and P.A. Maksym 648

Intrinsic bistability in nonlinear transport through a submicron lateral barrier
G. Pilling, D.H. Cobden, P.L. McEuen, C.I. Duruöz and J.S. Harris, Jr. 652

Conductance instabilities in quantum point contacts
J.C. Smith, C. Berven, M.N. Wybourne and S.M. Goodnick 656

Photoconductivity measurements of the electron–phonon interaction in quantum wire structures
A.J. Naylor, K.R. Strickland, A.J. Kent and M. Henini 660

Microwave-induced resonant reflection and localization of ballistic electrons in quantum microchannels
L.Y. Gorelik, M. Jonson and R.I. Shekhter 664

Surface acoustic wave studies of quantum nanostructures
G.R. Nash, S.J. Bending, Y. Kershaw, K. Eberl, P. Grambow and K. von Klitzing 668

Magnetotransport of electrons in arrays of wires in $Si/Si_{0.7}Ge_{0.3}$ heterostructures
M. Holzmann, D. Többen, P. Baumgartner, G. Abstreiter, A. Kriele, H. Lorenz and F. Schäffler 673

Spectroscopy of quantum dots in the few electron limit
Z. Borsosfoldi, I.A. Larkin, A.R. Long, M. Rahman, M.C. Holland, J.M.R. Weaver, J.H. Davies and J.G. Williamson 677

Coulomb blockade and dot size in split-gate wire with a small mesa or a hole made by scanning tunneling microscopy
S. Yamada, T. Kikutani and M. Yamamoto 682

**IX. Chaos, noise and fluctuations**

Chaos in quantum dots
B.L. Altshuler, N. Taniguchi and B.D. Simons 687

Berry phase, conductance oscillations and phase breaking in mesoscopic transport
Yu.B. Lyanda-Geller 692

Evidence for periodic "scar" patterns in the wavefunctions of a chaotic quantum well
P.B. Wilkinson, T.M. Fromhold, L. Eaves, F.W. Sheard, N. Miura and T. Takamasu 696

Persistent currents in the ballistic regime
R.A. Jalabert, K. Richter and D. Ullmo 700

Experimental investigation of Pauli exclusion principle using noise in coherent electron transport
Ç. Kurdak, C.-J. Chen, D.C. Tsui, J.P. Lu, M. Shayegan, S. Parihar and S.A. Lyon 705

Weak localisation in ballistic cavities filled with antidot arrays
G. Lütjering, K. Richter, D. Weiss, J. Mao, R.H. Blick, K. von Klitzing and C.T. Foxon 709

Calculations of magneto-transport fluctuations in ballistic quantum dots
G. Edwards, A. Grincwajg and D.K. Ferry 714

Conductance fluctuations in nanostructures of doped CdTe and $Cd_{1-x}Mn_xTe$ epilayers
J. Jaroszyński, J. Wróbel, M. Sawicki, T. Skośkiewicz, G. Karczewski, T. Wojtowicz, J. Kossut, T. Dietl, E. Kamińska, E.Papis and A. Piotrowska 718

Shot noise in the mesoscopic 2D diffusive systems in the limit of strong electron–electron scattering
V.I. Kozub and A.M. Rudin 722

Temporally correlated transport and shot noise suppression in a ballistic quantum point contact
M. Reznikov, M. Heiblum, H. Shtrikman and D. Mahalu 726

Phase breaking in ballistic quantum dots: a correlation field analysis
J.P. Bird, K. Ishibashi, D.K. Ferry, R. Newbury, D.M. Olatana, Y. Ochiai, Y. Aoyagi and T. Sugano 730

Multifractal statistics of eigenstates of 2D disordered conductors
V.I. Fal'ko and K.B. Efetov 735

Transport in one-dimensional electron Sinai billiards
M.V. Budantsev, Z.D. Kvon, A.G. Pogosov, N.T. Moschegov, A.E. Plotnikov and A.I. Toropov 739

Stable orbit bifurcations in a quantum well in a tilted magnetic field
G.S. Boebinger, G. Müller, H. Mathur, L.N. Pfeiffer and K.W. West 742

Coherent properties of in-plane gated InGaAs/AlGaAs submicron rings
A.A. Bykov, L.V. Litvin and S.P. Moshchenko 747

New *h/e* oscillations and electrostatic Aharonov–Bohm effect in a GaAs/AlGaAs-based mesoscopic ring
K. Park, S. Lee, M. Shin, E.-H. Lee and H.C. Kwon 751

**X. Quantum dots and wires – optical studies**

Optical spectroscopy of single quantum dots
A. Zrenner 756

Optical anisotropy of doped type I quantum wire arrays: from the zone center to the Fermi edge
T. Mélin and F. Laruelle 762

Coupled modes and filling factor dependent edge potentials in double-layered antidot arrays
K. Bollweg, T. Kurth, D. Heitmann, E. Vasiliadou, P. Grambow and K. Eberl 766

Exciton relaxation in self-organized InAs/GaAs quantum dots
R. Heitz, M. Grundmann, N.N. Ledentsov, L. Eckey, M. Veit, D. Bimberg, V.M. Ustinov, A.Yu. Egorov, A.E. Zhukov, P.S. Kop'ev and Zh.I. Alferov 770

Optical spectroscopies of electronic excitations in quantum dots
P. Hawrylak, A. Wojs, D.J. Lockwood, P.D. Wang, C.M. Sotomayor Torres, A. Pinczuk and B.S. Dennis 774

Temperature effects on the radiative recombination in self-assembled quantum dots
S. Fafard, S. Raymond, G. Wang, R. Leon, D. Leonard, S. Charbonneau, J.L. Merz, P.M. Petroff and J.E. Bowers 778

Resonant inelastic light scattering by electronic excitations in tunable quantum wire structures
G. Schedelbeck, E. Silveira, R. Strenz, U. Bockelmann, G. Abstreiter, G. Böhm and G. Weimann 783

Magneto-optical spectra in a doped 2-dimensional system with a periodic lateral modulation
M. Tavares and C. Tejedor 788

Magneto-optical studies of laterally modulated nanostructures grown on GaAs vicinal substrates
S. Huant and B. Etienne 793

Combined 1D–2D intersubband excitations and 1D single-particle spectrum in narrow quantum wires
G. Biese, C. Schüller, T. Kurth, D. Heitmann, P. Grambow and K. Eberl 797

Excitons in complex quantum nanostructures
G.W. Bryant, P.S. Julienne and Y.B. Band 801

Exciton interaction effects in the emission spectra of single free-standing InGaAs/GaAs quantum dots
R. Steffen, Th. Koch, J. Oshinowo, F. Faller and A. Forchel 805

Study of temperature-dependent exciton dynamics in a single quantum well with self-assembled islands
M.V. Marquezini, M.J.S.P. Brasil, J.A. Brum, P. Poole, S. Charbonneau and M.C. Tamargo 810

Naturally formed GaAs quantum dots
D. Gammon, E.S. Snow and D.S. Katzer 814

Effects of collective modes on single particle electronic excitations of semiconductor quantum wells and quantum dots
P.A. Knipp, S.W. Pierson and T.L. Reinecke 818

Magnetic field effects on the vacuum Rabi coupling in semiconductor quantum microcavity structures
T.A. Fisher, A.M. Afshar, D.M. Whittaker, M.S. Skolnick, J.S. Roberts, G. Hill and M.A. Pate 822

**XI. Electrons on helium**

Sliding of the Wigner solid on liquid helium
K. Kono and K. Shirahama 826

Evidence for quantum melting in the two-dimensional electron system on a thin helium film
T. Günzler, B. Bitnar, G. Mistura, S. Neser and P. Leiderer 831

Many-electron magnetoconductivity in 2D electrons on liquid helium
P.J. Richardson, A. Blackburn, K. Djerfi, M.I. Dykman, C. Fang-Yen, P. Fozooni, A. Kristensen, M.J. Lea and A. Santrich-Badal 835

The effect of confining electric fields on new magnetoplasma excitations in a 2D electron system
P.K.H. Sommerfeld, A.M.C. Valkering, R.W. van der Heijden and A.T.A.M. de Waele 839

Shear modes in 2D ion crystals trapped below the surface of superfluid helium
P.L. Elliott, A.A. Levchenko, C.I. Pakes, L. Skrbek and W.F. Vinen 843

**XII. Lateral superlattices and ballistics**

Interplay of Landau band width and band separation in a square lateral superlattice
T. Schlösser, K. Ensslin, J.P. Kotthaus and M. Holland 847

Magneto-oscillations of the gate current in a laterally modulated two-dimensional electron gas
F.A.P. Blom, F.M. Peeters, K. v.d. Zanden and M. van Hove 851

Magnetoresistance oscillations in a dimpled two-dimensional electron gas
G.M. Gusev, U. Gennser, X. Kleber, D.K. Maude, J.C. Portal, D.I. Lubyshev, P. Basmaji, M. de P.A. Silva, J.C. Rossi and Yu.V. Nastaushev 855

From low to high field: magneto-transport properties of electron gases in lateral superlattices grown on GaAs vicinal surfaces
L. Sfaxi, F. Petit, F. Lelarge, A. Cavanna and B. Etienne 860

**XIII. Novel techniques**

Quantum interference in 2D atomic-scale structures
M.F. Crommie, C.P. Lutz, D.M. Eigler and E.J. Heller 864

Nanofabrication with proximal probes
P.M. Campbell, E.S. Snow and P.J. McMarr 870

Fabrication of metallic nanostructures with an atomic force microscope
L.L. Sohn and R.L. Willett 874

Room temperature manipulation of $C_{60}$ molecules on a Si surface
P.H. Beton, A.W. Dunn and P. Moriarty 878

Quantum size effects in the optical conductivity of ultrathin metal films
L.A. Kuzik, V.A. Yakovlev, F.A. Pudonin and G. Mattei 882

Novel technique for fabrication of one- and two-dimensional systems
V.Ya. Prinz, V.A. Seleznev and A.K. Gutakovsky 886

Coulomb staircase with negative differential resistance at room temperature for a metal tip–metal dot–semiconductor double junction
P. Radojkovic, E. Hartmann, M. Schwartzkopff, M. Enachescu, E. Stefanov and F. Koch 890

**XIV. Organic conductors**

Magnetic-field dependent Fermi surfaces in quasi-2D organic conductors
J. Singleton, A.A. House, N. Harrison, I. Deckers, S.J. Blundell, J.A.A.J. Perenboom, A. Polisskii, W. Hayes, F. Herlach, M. Kurmoo and P. Day 894

Magneto-oscillations and field-induced phase transitions in organic conductors
R.H. McKenzie, J.S. Brooks, R.G. Clark, R. Newbury, R.P. Starrett, A. Skougarevsky, R.A. Lewis, S. Takasaki, J. Yamada, H. Anzai, Y. Tanaka, T. Kinoshita, N. Kinoshita, M. Tokumoto and N.D. Kushch 901

Novel tunneling experiments on organic heterostructures
C.M. Fischer, M. Burghard, S. Roth and K. von Klitzing 905

Comparative torque studies of $\alpha$-(BEDT-TTF)$_2$MHg(SCN)$_4$ (M = K, Tl, NH$_4$)
P. Christ, W. Biberacher, A.G.M. Jansen, M.V. Kartsovnik, A.E. Kovalev, N.D. Kushch, E. Steep and K. Andres 909

High-field Shubnikov–de Haas effect in the organic superconductor $\alpha$-(BEDT-TTF)$_2$NH$_4$Hg(SCN)$_4$
P.S. Sandhu, G.J. Athas, J.S. Brooks, E.G. Haanappel, J.D. Goettee, D.W. Rickel, M. Tokumoto, N. Kinoshita, T. Kinoshita and Y. Tanaka 913

**XV. Localisation and metal–insulator transition**

Inelastic transport in a system of 2D electrons localized in a smooth potential: giant negative magnetoresistance
M.E. Raikh and L.I. Glazman 918

Weak levitation of delocalized states as a result of Landau level mixing
T.V. Shahbazyan and M.E. Raikh 922

Metal–non-metal transition at the crossover from antidots to quantum dots
G. Lütjering, D. Weiss, R.W. Tank, K. von Klitzing, A. Hülsmann, T. Jakobus and K. Köhler 925

Semimetal–semiconductor transition in InAs/GaSb heterostructures
Jennifer J. Quinn and John J. Quinn 930

Magnetotransport anomalies in dilute two-dimensional electron systems: an experiment and a model
S.I. Dorozhkin, C.J. Emeleus, O.A. Mironov, T.E. Whall and G. Landwehr 933

Re-entrant metal–insulator transitions in Si–SiGe–Si heterostructures
M. D'Iorio, D. Stewart, S. Deblois, D. Brown and J.-P. Noël 937

Electric field induced non-ohmic conduction in the 2D insulating phase
V.M. Pudalov, M. D'Iorio and J.W. Campbell 941

The Coulomb gap and the transition to Mott hopping
F.W. van Keuls, X.L. Hu, A.J. Dahm and H.W. Jiang 945

Scaling of a metal/insulator transition in a 2D system at $B=0$
J.E. Furneaux, S.V. Kravchenko, W. Mason, V.M. Pudalov and M. D'Iorio 949

Experimental evidence of the Coulomb gap in a high-mobility 2D electron system in silicon
W. Mason, S.V. Kravchenko and J.E. Furneaux 953

Author index 957

Subject index 972

Materials index 989